

4

AD-A196 057

CCS Research Report 592

INTERACTING STRATEGY SETS IN
MULTIOBJECTIVE COMPETITION;
A DOMINANCE CONE CONSTRAINED
GAME SOLUTION


by

A. Charnes
Z.M. Huang
J.J. Rousseau
J. Semple

CENTER FOR CYBERNETIC STUDIES

The University of Texas
Austin, Texas 78712

SDTIC
ELECTE
JUL 13 1988
E



This document has been approved
for public release and sale in
distribution is unlimited.

88 7 13 05 9

4

CCS Research Report 592

INTERACTING STRATEGY SETS IN
MULTIOBJECTIVE COMPETITION;
A DOMINANCE CONE CONSTRAINED
GAME SOLUTION

by

A. Charnes
Z.M. Huang
J.J. Rousseau
J. Semple

April, 1988

The research for this paper was partly supported by National Science Foundation Grant SES 8520806 and U.S. Army Contract DAKF-15-87-C-0110, with the Center for Cybernetic Studies at The University of Texas at Austin. Reproduction in whole or in part is permitted for any purpose of the U.S. Government.

CENTER FOR CYBERNETIC STUDIES

A. Charnes, Director

College of Business Administration, 5.202
The University of Texas at Austin
Austin, Texas 78712-1177
(512) 471-1821

DTIC
SELECTED
JUL 13 1988
E

This document has been approved
for public release and sale in
unlimited quantities.

FOURTH INTERNATIONAL CONFERENCE ON THE FOUNDATIONS
AND APPLICATIONS OF UTILITY, RISK AND DECISION THEORY

ABSTRACT

Interacting Strategy Sets in Multiobjective
Competition; A Dominance Cone Constrained Game Solution

by

A. Charnes, Z. Huang, J.J. Rousseau, J. Semple
(The University of Texas at Austin)

Models employed in evaluation or stipulation of regulatory policies involving competition and/or differing objectives of competing parties have sometimes been seriously deficient in accounting for interactions between the parties' strategies or in allowing for multiple objectives. Classical game-theoretical models, wherein the strategy set is the topological product of the individual parties strategy sets, do not encompass such situations. The new "dominance cones" method and class of solutions (Charnes, Cooper, Huang, Wei) is herein further extended to such extensions of classical games. This is applied to an example of Harker without requiring his variational and quasi-variational inequalities or point-to-set mappings. *Key word: game theory.*


KEY WORDS

Dominance Cones

Multipayoff Cross-Constrained Games

Variational Inequalities

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



INTERACTING STRATEGY SETS IN MULTIOBJECTIVE COMPETITION; A DOMINANCE CONE CONSTRAINED GAME SOLUTION

by

A. Charnes, Z.M. Huang, J.J. Rousseau, and J. Semple

1. Introduction

Models employed in evaluation or stipulation of regulatory policies involving competition and/or differing objectives of competing parties have sometimes been seriously deficient in accounting for interactions between the parties' strategies or in allowing for multiple objectives. Classical game-theoretical models, wherein the strategy set is the topological product of the individual parties strategy sets, do not treat such situations. They can however be handled by the new "dominance cones" method and class of solutions (Charnes, Cooper, Wei and Huang [2]) to such extensions of classical games. Herein we extend the "C²WH" method to obtain more general results looking forward to resolution (in a later paper) of a competitive situation in transportation with interacting strategy sets [4].

Harker (1986) [5] used the Variational Inequality (VI) method to discuss the Generalized Nash Equilibrium Games (GNE) and gave an example in which only one solution can be found by (VI). Using the "dominance cones" method given in our paper, we can also find all the (GNE) solutions of the example given by Harker.

2. Nondominated Equilibrium Points

Definition 2.1: Let S be a set in E^m , the set $S^* = \{y \in E^m : x^T y \leq 0 \text{ for all } x \in S\}$ is called the negative polar cone of S .

Definition 2.2: Let Λ be a cone in E^m . Λ is said to be "acute" if there exists an open half-space

$$H = \{x \in E^m : a^T x > 0, a \neq 0\}$$

such that

$$\bar{\Lambda} \subset H \cup \{0\}$$

Lemma 2.1: Let Λ and Λ_1 be cones in E^m

- (i) If $\Lambda \subset \Lambda_1$, then $\Lambda^* \supset \Lambda_1^*$.

- (ii) $\text{Int } \Lambda^* \neq \emptyset$ if and only if Λ is acute.
- (iii) When Λ is acute, $\text{Int } \Lambda^* = \{y \in E^m : x^T y < 0 \text{ for all } x \in \bar{\Lambda} \text{ and } x \neq 0\}$ and $\bar{\Lambda} \cap (-\Lambda) = \{0\}$.
- (iv) If Λ is a convex cone, then $(\Lambda^*)^* = \bar{\Lambda}$.

Definition 2.3: Let S be a set in E^m , $\bar{x} \in \bar{S}$.

The tangency cone of S at \bar{x} is denoted by $T(S, \bar{x})$:

$$T(S, \bar{x}) = \{h \in E^m : \text{there exists a sequence } \{x^k\} \text{ and a sequence } \{\lambda^k\} \text{ such that}$$

$$h = \lim_{k \rightarrow \infty} \lambda_k (x^k - \bar{x}), \text{ where } x^k \in S, \lambda_k > 0 \text{ and } \lim_{k \rightarrow \infty} x_k = \bar{x}\}$$

For definitions and properties of cones, polar cones and direction cones the reader is referred to [1], [3], [6], [7], [8] and [9].

Definition 2.4: Let S be a convex set in E^m , Λ be a convex cone in E^n . A real-valued vector function

$G : S \rightarrow E^n$ is Λ -concave on S if

$$G(\lambda x^1 + (1-\lambda)x^2) - (\lambda G(x^1) + (1-\lambda)G(x^2)) \in \Lambda$$

for all $x^1, x^2 \in S$ and $\lambda \in (0,1)$.

Lemma 2.2: Let Λ be a closed convex cone in E^n , S be a convex set in E^m , $G : S \rightarrow E^n$ be

differentiable in a open set which contains S . If G is Λ -concave on S , then for every

$x^1, x^2 \in S$, we have

$$G(x^1) + \nabla_x G(x^1)(x^2 - x^1) \in G(x^2) + \Lambda$$

Proof: Since G is Λ -concave on S , for every $x^1, x^2 \in S$

$$G(\lambda x^2 + (1-\lambda)x^1) - (\lambda G(x^2) + (1-\lambda)G(x^1)) \in \Lambda \text{ for all } \lambda \in (0,1)$$

Thus

$$G(x^1) + \left[G(x^1 + \lambda(x^2 - x^1)) - G(x^1) \right] / \lambda \in G(x^2) + \Lambda$$

Letting $\lambda \rightarrow 0^+$, we have

$$G(x^1) + \nabla_x G(x^1)(x^2 - x^1) \in G(x^2) + \Lambda$$

Q.E.D.

Lemma 2.3: Let S be a convex set in E^m , Λ be a convex cone in E^n , $G : S \rightarrow E^n$. If G is Λ -concave on S , then for every $p \in (-\Lambda^*)$, $P^T G$ is concave on S .

Proof: Since G is Λ -concave on S , for all $x^1, x^2 \in S$ and all $\lambda \in (0,1)$, we have

$$G(\lambda x^1 + (1-\lambda)x^2) - (\lambda G(x^1) + (1-\lambda)G(x^2)) \in \Lambda$$

Thus for every $p \in (-\Lambda^*)$, we have

$$p^T G(\lambda x^1 + (1-\lambda)x^2) \geq \lambda p^T G(x^1) + (1-\lambda)p^T G(x^2)$$

for all $\lambda \in (0,1)$.

Q.E.D.

Lemma 2.4: Let S be a convex set in E^m , Λ be a closed convex cone in E^n , $G : S \rightarrow E^n$. If, for arbitrary $p \in \Lambda^*$, $P^T G$ is concave on S , then G is $(-\Lambda)$ -concave on S .

Proof: Since for arbitrary $p \in \Lambda^*$ and $x, y \in S$, we have

$$p^T G(\lambda x + (1-\lambda)y) \geq \lambda p^T G(x) + (1-\lambda)p^T G(y) \text{ for all } \lambda \in (0,1)$$

i.e.,

$$p^T \{G(\lambda x + (1-\lambda)y) - (\lambda G(x) + (1-\lambda)G(y))\} \geq 0 \text{ for all } \lambda \in (0,1)$$

By Lemma 2.1,

$$G(\lambda x + (1-\lambda)y) - (\lambda G(x) + (1-\lambda)G(y)) \in -(\Lambda^*)^* = -\Lambda$$

for all $\lambda \in (0,1)$.

This means that G is $(-\Lambda)$ -concave on S .

Q.E.D.

A Multi-Payoff Constrained N-person Game In Normal Form

Definition 2.5: A multi-payoff constrained n -person game in normal form is given by n nonempty sets S_i in E^n ($i = 1, 2, \dots, n$), the strategy sets of the players $1, 2, \dots, n$; a real-valued vector function $G = (g_1, \dots, g_m)^T : S_1 \times \dots \times S_n \rightarrow E^m$, the cross-constraint function of the n players; for each player i there is a real-valued vector function $A^i = (A_1^i, \dots, A_l^i)^T : S_1 \times \dots \times S_n \rightarrow E^l$, the vector payoff function of the i th player ($i = 1, 2, \dots, n$); a convex cone K in E^m , the constraint cone; a convex cone W in E^l , the dominance cone; $\mathfrak{R}_K = \{x = (x_1, \dots, x_n) : G(x) \in K, x_i \in S_i, i = 1, 2, \dots, n\}$, the constraint set.

Such a game will be denoted by

$$\Gamma = \{ \mathfrak{R}_K, W; A^1, \dots, A^n \}.$$

Definition 2.6: Let $\Gamma = \{ \mathfrak{R}_K, W; A^1, \dots, A^n \}$ be a multi-payoff constrained n-person game.

$\bar{x} = (\bar{x}_1, \dots, \bar{x}_n) \in \mathfrak{R}_K$ is called a nondominated equilibrium point of the game Γ associated with

W if there exists no $x = (x_1, \dots, x_n) \in S_1 \times \dots \times S_n$ satisfying

$(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \in \mathfrak{R}_K$ ($i = 1, 2, \dots, n$) such that

$$A^i(\bar{x}_1, \dots, \bar{x}_n) \in A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) + W$$

and

$$A^i(\bar{x}_1, \dots, \bar{x}_n) \neq A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n)$$

for all $i = 1, \dots, n$.

Definition 2.7: Let $\Gamma = \{ \mathfrak{R}_K, W; A^1, \dots, A^n \}$ be a multi-payoff constrained n-person game. Then Γ is

called a (W-K)-concave game if the following four conditions hold for all $i = 1, 2, \dots, n$:

- (i) S_i is a convex set in E^{n_i} ;
- (ii) $A^i(x_1, \dots, x_n)$ is (-W)-concave with respect to x_i on S_i for fixed $x_j \in S_j$ ($j \neq i, j = 1, 2, \dots, n$);
- (iii) $A^i(x_1, \dots, x_n)$ is continuous on $S_1 \times \dots \times S_n$.
- (iv) $G(x_1, \dots, x_n)$ is continuous and K-concave on $S_1 \times \dots \times S_n$.

In the remaining sections, except where specifically noted, we shall always use the following

symbols:

$$(1) \quad W_n = \underbrace{W \times \dots \times W}_n, \quad W_n^* = \underbrace{W^* \times \dots \times W^*}_n, \quad K_n = \underbrace{K \times \dots \times K}_n$$

$$K_n^* = K^* \times \dots \times K^* \text{ and } S = S_1 \times \dots \times S_n;$$

$$(2) \quad \text{"Strictly nonzero"} \quad p \in W_n^* \text{ implies } p = (p^1, \dots, p^n), \quad p^i \in W^* \text{ and } p^i \neq 0 \text{ for all}$$

$$i = 1, 2, \dots, n;$$

$$(3) \quad Q \in -K_n^* \text{ implies } Q = (q^1, \dots, q^n), \quad q^i \in -K^*, \quad i = 1, 2, \dots, n;$$

$$(4) \quad x \in S \text{ implies } x = (x_1, \dots, x_n) \text{ and } x_i \in S_i \text{ for all } i = 1, 2, \dots, n;$$

- (5) For fixed $x \in \mathfrak{R}_K$, let for all $i = 1, 2, \dots, n$

$$D_i(\bar{x}) = \left\{ x_i \in S_i : (\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \in \mathfrak{R}_K \right\}$$

and let $D(\bar{x}) \triangleq D_1(\bar{x}) \times \dots \times D_n(\bar{x})$.

Clearly, for all $i = 1, 2, \dots, n$, we have $D_i(\bar{x}) \subset S_i$ and $D(\bar{x}) \subset S$.

- (6) $x \in D(\bar{x})$ implies for $x = (x_1, \dots, x_n)$ that $x_i \in D_i(\bar{x})$ for all $i = 1, 2, \dots, n$.
- (7) For a real-valued vector function $G(x) = (g_1(x), \dots, g_m(x))^T$, we denote the "gradient" of $G(x)$ (really, the vector of gradients of G 's n component) by

$$\nabla_{x_i} G(x) = \begin{pmatrix} \nabla_{x_i} g_1(x) \\ \vdots \\ \nabla_{x_i} g_m(x) \end{pmatrix}, \quad (x_i \text{ is a vector!})$$

- (8) For a specific $p \in W_n^*$, let

$$\Psi_p(x, y) = \sum_{i=1}^n p^i A^i(y_1, \dots, y_{i-1}, x_i, y_{i+1}, \dots, y_n) \text{ for all } (x, y) \in S \times S.$$

- (9) For a specific $p \in W_n^*$, the generalized Lagrangean function is defined as follows:

$$\Phi_p(x, y, Q) = \sum_{i=1}^n p^i A^i(y_1, \dots, y_{i-1}, x_i, y_{i+1}, \dots, y_n) + \sum_{i=1}^n q^i G(y_1, \dots, y_{i-1}, x_i, y_{i+1}, \dots, y_n)$$

for all $(x, y) \in S \times S$ and $Q \in -K_n^*$.

3. Computing Nondominated Equilibrium Points

In this section we outline a more general technique than was given previously in [2], which enables us to capture additional nondominated equilibrium points.

Lemma 3.1: Let $\Gamma = \{\mathfrak{R}_K, W; A^1, \dots, A^n\}$ be a $(W-K)$ -concave game. If $\bar{x} \in \mathfrak{R}_K$ is a nondominated equilibrium point of Γ associated with W , then there exists strictly nonzero $p \in W_n^*$ such that for arbitrary $x \in D(\bar{x})$, we have

$$p^i A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \leq p^i A^i(\bar{x}) \text{ for all } i = 1, 2, \dots, n.$$

Proof: Suppose that $\bar{x} \in \mathfrak{R}_K$ is a nondominated equilibrium point of Γ associated with W . This means that there exist no n nonzero $w_i \in W$ ($i = 1, 2, \dots, n$) and $x \in D(\bar{x})$ such that

$$A^i(\bar{x}) = A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) + w_i \text{ for all } i = 1, 2, \dots, n.$$

For each $i = 1, 2, \dots, n$, consider

$$\Lambda_i = \left\{ z \in E^1 : z - A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) + A^i(\bar{x}) = w_i \right. \\ \left. \text{for some } x_i \in D_i(\bar{x}) \text{ and nonzero } w_i \in W \right\}$$

It is easy to show that Λ_i is a convex set and $0 \notin \Lambda_i$. Hence by the separation theorem, there exists nonzero $p^i \in E^1$ such that

$$x p^{i^T} z \leq 0 \quad \text{for all } z \in \Lambda_i.$$

Next, for arbitrary $x_i \in D_i(\bar{x})$, nonzero $w_i \in W$ and $\lambda > 0$, let

$$z_{x_i \lambda w_i} = A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) - A^i(\bar{x}) + \lambda w_i.$$

Then $z_{x_i \lambda w_i} \in \Lambda_i$ and

$$p^{i^T} A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) - p^{i^T} A^i(\bar{x}) + \lambda p^{i^T} w_i \leq 0.$$

Thus

$p^i \in W^*$. Letting $\lambda \rightarrow 0^+$, we obtain

$$p^{i^T} A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \leq p^{i^T} A^i(\bar{x}) \text{ for all } i = 1, 2, \dots, n.$$

Q.E.D.

Lemma 3.2: Let $\Gamma = \{\mathfrak{R}_K, W; A^1, \dots, A^n\}$, $\bar{x} \in \mathfrak{R}_K$ and $p \in \text{Int } W_n^*$ if for arbitrary $x \in D(\bar{x})$, we have

$$p^{i^T} A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \leq p^{i^T} A^i(\bar{x}) \quad (3.1)$$

for all $i = 1, 2, \dots, n$, then \bar{x} is a nondominated equilibrium point of Γ associated with W .

Proof: Suppose to the contrary that \bar{x} is not a nondominated equilibrium point of Γ associated with W . That is, there exist $x \in D(\bar{x})$ and $w_i \in W$ such that

$$A^i(\bar{x}) = A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) + w_i, \quad w_i \neq 0$$

for all $i = 1, 2, \dots, n$.

We note by Lemma 2.1 that W is an acute cone. By the acute property of W and Lemma 2.1, we have $p^i w_i < 0$, hence

$$p^i A^i(\bar{x}) < p^i A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \text{ for all } i = 1, \dots, n.$$

This contradicts (3.1).

Q.E.D.

Lemma 3.3: Let $\Gamma = \{\mathcal{R}_K, W; A^1, \dots, A^n\}$ be a (W-K)-concave game. If $\bar{x} \in \mathcal{R}_K$ is a nondominated equilibrium point of Γ associated with W , then there exists strictly nonzero $\bar{p} \in W_n^+$ such that

$$\Psi_{\bar{p}}(x, \bar{x}) \leq \Psi_{\bar{p}}(\bar{x}, \bar{x}) \quad \text{for all } x \in D(\bar{x}).$$

Proof: By Lemma 3.1, there exists a strictly nonzero $\bar{p} \in W_n^+$ such that for arbitrary $x \in D(\bar{x})$, we have

$$\bar{p}^i A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \leq \bar{p}^i A^i(\bar{x})$$

for all $i = 1, 2, \dots, n$, then

$$\sum \bar{p}^i A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \leq \sum_{i=1}^n \bar{p}^i A^i(\bar{x})$$

$$\text{i.e.,} \quad \Psi_{\bar{p}}(x, \bar{x}) \leq \Psi_{\bar{p}}(\bar{x}, \bar{x}) \quad \text{for all } x \in D(\bar{x})$$

Q.E.D.

Lemma 3.4: Let $\Gamma = \{\mathcal{R}_K, W; A^1, \dots, A^n\}$, $\bar{x} \in \mathcal{R}_K$ and $A^i(x)$ be differentiable at \bar{x} with respect to x_i for fixed $\bar{x}_1, \dots, \bar{x}_{i-1}, \bar{x}_{i+1}, \dots, \bar{x}_n$ ($i = 1, \dots, n$). If there exists $\bar{p} \in W_n^+$ such that

$$\Psi_{\bar{p}}(x, \bar{x}) \leq \Psi_{\bar{p}}(\bar{x}, \bar{x}) \quad \text{for all } x \in D(\bar{x})$$

then

$$\left(\nabla_{x_i} \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} \right)^T \in T^*(D_i(\bar{x}), \bar{x}_i) \quad (i = 1, \dots, n)$$

Proof: For arbitrary $h \in T(D_i(\bar{x}), \bar{x}_i)$, there exists $\{x_i^k\} \subset D_i(\bar{x})$ with

$$\lim_{k \rightarrow \infty} x_i^k = \bar{x}_i, \text{ and } \lambda_k > 0 \text{ with } \lambda_k \rightarrow 0 \text{ such that}$$

$$h = \lim_{k \rightarrow \infty} \lambda_k (x_i^k - \bar{x}_i)$$

since $\Psi_{\bar{p}}(x, \bar{x})$ is differentiable at \bar{x} , we have

$$\Psi_{\bar{p}}(x^k, \bar{x}) = \Psi_{\bar{p}}(\bar{x}, \bar{x}) + \nabla_{x_i} \Psi_{\bar{p}}(x^k, \bar{x}) \Big|_{x=\bar{x}} (x_i^k - \bar{x}_i) + \|x_i^k - \bar{x}_i\| \cdot \varepsilon(k)$$

where $x^k = (\bar{x}_1, \dots, \bar{x}_{i-1}, x_i^k, \bar{x}_{i+1}, \dots, \bar{x}_n)$, $\varepsilon(k) \rightarrow 0$ as $k \rightarrow \infty$.

Since $\Psi_{\bar{p}}(\bar{x}, \bar{x}) \geq \Psi_{\bar{p}}(x, \bar{x})$ for all $x \in \mathcal{R}_K$, we have

$$\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} (x_i^k - \bar{x}_i) + \left\| x_i^k - \bar{x}_i \right\| \bullet \varepsilon(k) \leq 0$$

then

$$\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} \lambda_k (x_i^k - \bar{x}_i) + \left\| \lambda_k (x_i^k - \bar{x}_i) \right\| \bullet \varepsilon(k) \leq 0$$

Letting $k \rightarrow \infty$, we have

$$\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} h \leq 0$$

Hence

$$\left(\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} \right)^T \in T^*(D_i(\bar{x}), \bar{x}_i)$$

Q.E.D

Lemma 3.5. Let $\Gamma = \{\mathcal{R}_K, W; A^1, \dots, A^n\}$ be a (W-K)-concave game and $\bar{x} \in \mathcal{R}_K$. For all $i = 1, 2, \dots, n$, let $A^i(x)$ be differentiable at \bar{x} with respect to x_i for fixed $\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n$ ($i = 1, \dots, n$). If there exists $\bar{p} \in W_n$ such that \bar{x} satisfying

$$\left(\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} \right) (y - \bar{x}) \leq 0 \text{ for all } y \in D(\bar{x}),$$

then \bar{x} is a nondominated equilibrium point of Γ associated with W .

Proof:

By Lemma 3.2, we only need to show that for all $x \in D(\bar{x})$, we have

$$\bar{p}^T A^i(\bar{x}_1, \dots, \bar{x}_{i-1}, x_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \leq \bar{p}^T A^i(\bar{x}), \quad i = 1, \dots, n.$$

Suppose to the contrary that there exist $\bar{y} \in D(\bar{x})$ and some i_0 ($1 \leq i_0 \leq n$) such that

$$\bar{p}^{i_0 T} A^{i_0}(\bar{x}_1, \dots, \bar{x}_{i_0-1}, x_{i_0}, \bar{x}_{i_0+1}, \dots, \bar{x}_n) > \bar{p}^{i_0 T} A^{i_0}(\bar{x}). \quad (3.2)$$

Since Γ is a (W-K)-concave game, by Lemma 2.3, $\bar{p}^{i_0 T} A^{i_0}(\bar{x}_1, \dots, \bar{x}_{i_0-1}, x_{i_0}, \bar{x}_{i_0+1}, \dots, \bar{x}_n)$ is concave

with respect to x_{i_0} and thus

$$\bar{p}^{i_0 T} A^{i_0}(\bar{x}_1, \dots, \bar{x}_{i_0-1}, x_{i_0}, \bar{x}_{i_0+1}, \dots, \bar{x}_n) \leq \bar{p}^{i_0 T} A^{i_0}(\bar{x}) + \bar{p}^{i_0 T} \nabla_{x_{i_0}} A^{i_0}(\bar{x}) (\bar{y}_{i_0} - \bar{x}_{i_0}) \quad (3.3)$$

Since

$$\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} (y - \bar{x}) \leq 0 \text{ for all } y \in D(\bar{x}),$$

Letting $y = (\bar{x}_1, \dots, \bar{x}_{i_0-1}, x_{i_0}, \bar{x}_{i_0+1}, \dots, \bar{x}_n)$, we have

$$\nabla_{x_{i_0}} \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} (\bar{y}_{i_0} - \bar{x}_{i_0}) \leq 0$$

$$\text{i.e.,} \quad \bar{p}^{i_0} \nabla_{x_{i_0}} A^{i_0}(\bar{x}) (\bar{y}_{i_0} - \bar{x}_{i_0}) \leq 0$$

By (3.3), we have

$$\bar{p}^{i_0} A^{i_0}(\bar{x}_1, \dots, \bar{x}_{i_0-1}, x_{i_0}, \bar{x}_{i_0+1}, \dots, \bar{x}_n) \leq \bar{p}^{i_0} A^{i_0}(\bar{x})$$

This contradicts (3.2).

Q.E.D

Assuming $\bar{x} \in \mathfrak{R}_K$ let

$$c_i(\bar{x}) = \left\{ \left(\nabla_{x_i} G(\bar{x}) \right)^T : q \in (-K^*) \text{ such that } q^T G(\bar{x}) = 0 \right\}$$

Lemma 3.6^[1] Let $\bar{x} \in \mathfrak{R}_K$ and $G(x)$ be differentiable at \bar{x} .

Then

$$T(D_i(\bar{x}), \bar{x}_i) \subset (-C^*, i(\bar{x}))$$

Definition 3.1: A point $\bar{x} \in \mathfrak{R}_K$ is said to be a "generalized regular point" of the constraint set \mathfrak{R}_K if

$$T^*(D(\bar{x}), \bar{x}) \subset (-C^*, i(\bar{x})).$$

Definition 3.2: If there exist

$$\bar{x} = \bar{x}_1, \dots, \bar{x}_n \in E^{n_1} \times \dots \times E^{n_n}, \bar{Q} \in -K^*_{-n} \text{ and strictly nonzero } \bar{p} \in W^*_{-n} \text{ such that}$$

$$\nabla_{x_i} \Phi_{\bar{p}}(x, \bar{x}, \bar{Q}) \Big|_{x=\bar{x}} = \bar{p}^{i^T} \nabla_{x_i} A^i(\bar{x}) + \bar{q}^{i^T} \nabla_{x_i} G(\bar{x}) = 0, \bar{x}_i \in S_i \quad (3.4)$$

$$\nabla_{q_i} \Phi_{\bar{p}}(x, \bar{x}, \bar{Q}) = G(\bar{x}) \in K \quad (3.5)$$

$$\bar{q}^{i^T} \nabla_{q_i} \Phi_{\bar{p}}(x, \bar{x}, \bar{Q}) = \bar{q}^{i^T} G(\bar{x}) = 0 \quad (3.6)$$

for all $i = 1, \dots, n$, then we say that $(\bar{x}, \bar{p}, \bar{Q})$ satisfies the "generalized Kuhn Tucker" conditions.

Theorem 3.1: Let $\Gamma = \{\mathfrak{R}_K, W; A^1, \dots, A^n\}$ be a (W-K)-concave game. Furthermore, suppose that

$\bar{x} \in \mathfrak{R}_K$ is a nondominated equilibrium point of Γ associated with W , and that \bar{x} is a "generalized regular point" of \mathfrak{R}_K . Then there exist $\bar{Q} \in -K^*_{-n}$ and strictly nonzero $\bar{p} \in W^*_{-n}$ such that

$(\bar{x}, \bar{p}, \bar{Q})$ satisfies the generalized Kuhn - Tucker conditions (3.4) - (3.6).

Proof: By Lemma 3.3, there exists a strictly nonzero $\bar{p} \in W^*_{-n}$ such that

$$\Psi_{\bar{p}}(x, \bar{x}) \leq \Psi_{\bar{p}}(\bar{x}, \bar{x}) \quad \text{for all } x \in D(\bar{x})$$

Since \bar{x} is a generalized regular point of \mathfrak{R}_K and

$$\left(\nabla_{x_i} \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} \right)^T \in T^*(D(\bar{x}), \bar{x}) \quad (\text{see Lemma 3.4}),$$

there exist $\bar{q}^i \in E^m$ such that

$$\nabla_{x_i} \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} + \bar{q}^i{}^T \nabla_{x_i} G(\bar{x}) = 0$$

and

$$\bar{q}^i \in (-K^*), \quad \bar{q}^i{}^T G(\bar{x}) = 0$$

for all $i = 1, 2, \dots, n$.

i.e.,

$$\nabla_{x_i} \Phi_{\bar{p}}(x, \bar{x}, \bar{Q}) \Big|_{x=\bar{x}} = \bar{p}^i{}^T \nabla_{x_i} A^i(\bar{x}) + \bar{q}^i{}^T \nabla_{x_i} G(\bar{x}) = 0, \quad \bar{x}_i \in S_i$$

$$\nabla_{q_i} \Phi_{\bar{p}}(\bar{x}, \bar{x}, \bar{Q}) = G(\bar{x}) \in K$$

$$\bar{q}^i{}^T \nabla_{q_i} \Phi_{\bar{p}}(\bar{x}, \bar{x}, \bar{Q}) = \bar{q}^i{}^T G(\bar{x}) = 0$$

for all $i = 1, 2, \dots, n$ where $\bar{Q} = (\bar{q}^1, \dots, \bar{q}^n) \in -K_n^*$

Q.E.D.

Theorem 3.2: Let $\Gamma = \{\mathfrak{R}_K, W; A^1, \dots, A^n\}$ be a (W-K)-concave game, let K be closed and $\bar{x} \in \mathfrak{R}_K$,

let A^i be differentiable at \bar{x} with respect to x_i for fixed $\bar{x}_1, \dots, \bar{x}_{i-1}, \bar{x}_{i+1}, \dots, \bar{x}_n$ and $G(x)$ be

differentiable at \bar{x} . If there exist $\bar{p} \in \text{Int } W_n^*$ and $\bar{Q} \in -K_n^*$ such that $(\bar{x}, \bar{p}, \bar{Q})$ satisfies the

generalized Kuhn-Tucker conditions (3.4) - (3.6), then \bar{x} is a nondominated equilibrium point of

Γ associated with W .

Proof: According to Lemma 3.5, it suffices to show that

$$\nabla_x \Psi_{\bar{p}}(x, \bar{x}) \Big|_{x=\bar{x}} (y - \bar{x}) \leq 0 \quad \text{for all } y \in D(\bar{x}),$$

For arbitrary $y \in \mathfrak{R}_K$, we have by Lemma 2.2,

$$G(\bar{x}) + \nabla_x G(\bar{x})(y - \bar{x}) \in G(y) + K$$

Hence

$$\bar{q}^i{}^T \Delta_{x_i} G(\bar{x})(y - \bar{x}) \geq 0 \quad \text{for any } y \in \mathfrak{R}_K$$

letting $y = (\bar{x}_1, \dots, \bar{x}_{i-1}, y_i, \bar{x}_{i+1}, \dots, \bar{x}_n) \in \mathfrak{R}_K$, then

$$\bar{q}^i{}^T \Delta_{x_i} G(\bar{x})(y - \bar{x}) \geq 0 \quad \text{for all } y_i \in D_i(\bar{x})$$

and then

$$\nabla_{x_i} \Psi_p(x, \bar{x}) \Big|_{x=\bar{x}} (y_i - \bar{x}_i) = q^i \nabla_{x_i} G(\bar{x}) (y_i - \bar{x}_i) \leq 0$$

for all $y_i \in D_i(\bar{x})$, $i = 1, 2, \dots, n$;

i.e.,

$$\nabla_{x_i} \Psi_p(x, \bar{x}) \Big|_{x=\bar{x}} (y - \bar{x}) \leq 0 \quad \text{for all } y \in D(\bar{x})$$

Q.E.D.

Now, let's consider the example in Harker's paper, the two -person game depicted in Fig. 1. Each player chooses a number x_i between 0 and 10, such that the sum of these numbers is less than or equal to 15. Harker's utility functions and constraint functions are defined in our terms by:

$$A_1(x_1, x_2) = 34x_1 - x_1^2 - (8/3)x_1x_2$$

$$A_2(x_1, x_2) = 24.25x_2 - x_2^2 - (5/4)x_1x_2$$

$$G(x_1, x_2) = (10 - x_1, x_1, 10 - x_2, x_2, 15 - x_1 - x_2)^T$$

Here $W = E_-^1$, $K = E_+^5$, $S = E^2$, $\mathcal{R}_K = \{x = (x_1, x_2)^T : G(x_1, x_2) \in K\}$

since $G(x)$ is linear, for any point $x \in \mathcal{R}_K$, x is a generalized regular point of \mathcal{R}_K . It is easy to check that $\Gamma = \{\mathcal{R}_K, W; A_1, A_2\}$ is a $(W - K)$ -concave game. Hence by Theorem 3.1 and Theorem 3.2, these nondominated equilibrium points of Γ are points which satisfy the generalized Kuhn-Tucker conditions (3.4) - (3.6) and vice-versa.

Let $\Phi(x, y, \lambda) = A_1(x_1, y_2) + A_2(y_1, x_2) + \lambda^1 G(x_1, y_2) + \lambda^2 G(y_1, x_2)$

where $\lambda = (\lambda^1, \lambda^2)$, $\lambda^1 = (\lambda_1^1, \dots, \lambda_5^1)^T$, $\lambda^2 = (\lambda_1^2, \dots, \lambda_5^2)^T$.

Then the generalized Kuhn-Tucker conditions of Γ are to find $(\bar{x}, \bar{\lambda}^1, \bar{\lambda}^2)$ satisfying:

$$\nabla_{x_1} \Phi(x, \bar{x}, \bar{\lambda}) \Big|_{x=\bar{x}} = 34 - 2\bar{x}_1 - (8/3)\bar{x}_2 - \bar{\lambda}_1^1 + \bar{\lambda}_2^1 - \bar{\lambda}_5^1 = 0$$

$$\nabla_{x_2} \Phi(x, \bar{x}, \bar{\lambda}) \Big|_{x=\bar{x}} = 24.25 - 2\bar{x}_2 - (5/4)\bar{x}_1 - \bar{\lambda}_3^2 + \bar{\lambda}_4^2 - \bar{\lambda}_5^2 = 0$$

$$\nabla_{\lambda_i} \Phi(\bar{x}, \bar{x}, \bar{\lambda}) = (10 - \bar{x}_1, \bar{x}_1, 10 - \bar{x}_2, \bar{x}_2, 15 - \bar{x}_1 - \bar{x}_2)^T \geq 0, \lambda^i \geq 0, i = 1, 2.$$

$$\bar{\lambda}^1 \nabla_{\lambda^1} \Phi(\bar{x}, \bar{x}, \bar{\lambda}) = \bar{\lambda}_1^1(10 - \bar{x}_1) + \bar{\lambda}_2^1 \bar{x}_1 + \bar{\lambda}_3^1(10 - \bar{x}_2) + \bar{\lambda}_4^1 \bar{x}_2 + \bar{\lambda}_5^1(15 - \bar{x}_1 - \bar{x}_2) = 0$$

Simplifying the above, this is equivalent to finding $(\bar{x}, \bar{\lambda}^1, \bar{\lambda}^2)$ satisfying the following system:

$$34 - 2x_1 - (8/3)x_2 = \lambda_1^1 - \lambda_2^1 + \lambda_5^1 \quad (3.7)$$

$$24.25 - 2x_2 - (5/4)x_1 = \lambda_3^2 - \lambda_4^2 + \lambda_5^2 \quad (3.8)$$

$$\lambda_1^i (10 - x_1) = 0, \quad x_1 \leq 10, \lambda_1^i \geq 0, \quad i = 1, 2 \quad (3.9)$$

$$\lambda_2^i x_1 = 0, \quad x_1 \geq 0, \lambda_2^i \geq 0, \quad i = 1, 2 \quad (3.10)$$

$$\lambda_3^i (10 - x_2) = 0, \quad x_2 \leq 10, \lambda_3^i \geq 0, \quad i = 1, 2 \quad (3.11)$$

$$\lambda_4^i x_2 = 0, \quad x_2 \geq 0, \lambda_4^i \geq 0, \quad i = 1, 2 \quad (3.12)$$

$$\lambda_5^i (15 - x_1 - x_2) = 0, \quad \lambda_5^i \geq 0, \quad x_1 + x_2 \leq 15 \quad (3.13)$$

The constraint set \mathcal{R}_K is illustrated in Fig. 1.




Fig. 1.

We divide \mathcal{R}_K into six parts I_i , $i = 0, 1, 2, 3, 4, 5$:

$$I_0 = \text{Int } \mathcal{R}_K = \{(x_1, x_2) : 0 < x_1 < 10, 0 < x_2 < 10, x_1 + x_2 < 15\}$$

$$I_1 = \{(x_1, x_2) : 0 \leq x_1 \leq 5, x_2 = 10\}$$

$$I_2 = \{(x_1, x_2) : 5 < x_1 \leq 10, x_1 + x_2 = 15\}$$

$$I_3 = \{(x_1, x_2) : x_1 = 0, 0 < x_2 < 5\}$$

$$I_4 = \{(x_1, x_2) : 0 < x_1 \leq 10, x_2 = 0\}$$

$$I_5 = \{(x_1, x_2) : x_1 = 0, 0 \leq x_2 < 10\}$$

For each part I_i , using the complementary conditions (3.9) - (3.13), it is easy to figure out which points satisfy the system (3.7) - (3.13). The process is the following:

(1) I_0 : Since $0 < x_1 < 10$, $0 < x_2 < 10$ and $x_1 + x_2 < 15$, by (3.9) - (3.13) we have $\lambda_j^i = 0$ $i = 1, 2$,

$j = 1, 2, 3, 4, 5$. By (3.7) and (3.8), we have

$$34 - 2x_1 - (8/3)x_2 = 0$$

$$24.25 - 2x_2 - (5/4)x_1 = 0$$

then $(\bar{x}_1, \bar{x}_2) = (5.9)$ satisfies (3.7) - (3.13)

(2) I_1 : Since $0 \leq x_1 \leq 5$ and $x_2 = 10$, by (3.9) and (3.12) we have $\lambda_1^i = 0$, $\lambda_4^i = 0$.

By (3.7) and (3.8), we have

$$34 - 2x_1 - (80/3) = -\lambda_2^1 + \lambda_5^1 \quad (3.14)$$

$$24.25 - 20 - (5/4)x_1 = \lambda_3^2 + \lambda_5^2 \quad (3.15)$$

(a) $x_1 = 0$. By (3.13) we have $\lambda_5^i = 0$, $i = 1, 2$.

By (3.14), $\lambda_2^1 = -22/3 < 0$, which contradicts (3.10).

(b) $x_1 = 5$. by (3.10) we have $\lambda_2^i = 0$, $i = 1, 2$.

By (3.14), $\lambda_5^1 = -8/3 < 0$, which contradicts (3.13).

(c) $0 < x_1 < 5$. By (3.10) and (3.13) we have $\lambda_2^i = \lambda_5^i = 0$ ($i = 1, 2$) and then by (3.14)

and (3.15), we have

$$x_1 = 11/3$$

$$\lambda_3^2 = -1/3 < 0, \text{ which contradicts (3.11).}$$

(3) I_2 : Since $5 < x_1 \leq 10$ and $x_1 + x_2 = 15$ (obviously $0 < x_2 = 15 - x_1 < 10$), by (3.10) - (3.12) we have

$$\lambda_2^i = \lambda_3^i = \lambda_4^i = 0, \quad i = 1, 2.$$

then by (3.7) and (3.8), we have

$$34 - 2x_1 - (8/3)(15 - x_1) = \lambda_1^1 + \lambda_5^1$$

$$24.25 - 2(15 - x_1) - (5/4)x_1 = \lambda_5^2$$

i.e.,

$$\lambda_5^1 = (2/3)x_1 - 6 - \lambda_1^1 \quad (3.16)$$

$$\lambda_5^2 = (3/4)x_1 - 5.75 \quad (3.17)$$

(a) $x_1 = 10$. By (3.16) and (3.17), we have

$$\lambda_1^1 + \lambda_5^1 = 2/3 \quad (3.18)$$

$$\lambda_5^2 = 7/4 \quad (3.19)$$

so we can find $\lambda_1^1 \geq 0$, $\lambda_5^1 \geq 0$, $\lambda_5^2 = 7/4$ to satisfy (3.18) and (3.19),

i.e., (10, 5) satisfies the generalized Kuhn - Tucker conditions.

(b) $x_1 < 10$. By (3.9) we have $\lambda_i^1 = 0$ $i = 1, 2$. Then by (3.16) and (3.17), we have

$$\lambda_5^1 = (2/3)x_1 - 6$$

$$\lambda_5^2 = (3/4)x_1 - 5.75$$

letting $\lambda_5^1 \geq 0$, $\lambda_5^2 \geq 0$, we have

$$x_1 \geq 9 \text{ and } x_1 \geq 23/3$$

but $x_1 \geq 9$ implies $x_1 \geq 23/3$.

Combining (a) and (b), the segment

$$I_1^2 = \{(x_1, x_2) : 9 \leq x_1 \leq 10, x_2 = 15 - x_1\}$$

is the solution set of system (3.7) - (3.13) in I_2 , i.e., that satisfying the generalized Kuhn - Tucker conditions.

(4) I_3 : Since $x_1 = 10$, $0 < x_2 < 5$, by (3.10) - (3.13) we have

$$\lambda_2^i = \lambda_3^i = \lambda_4^i = \lambda_5^i = 0, \quad i = 1, 2$$

By (3.7) and (3.8), we have

$$34 - 20 - (8/3)x_2 = \lambda_1^1$$

$$24.25 - 2x_2 - 50/4 = 0$$

then

$$x_2 = 47/8$$

$$\lambda_1^1 = -5/3 < 0, \text{ which contradicts (3.9).}$$

(5) I_4 : Since $0 < x_1 \leq 10$, $x_2 = 0$, by (3.10), (3.11) and (3.13) we have

$$\lambda_2^i = \lambda_3^i = \lambda_5^i = 0, \quad i = 1, 2$$

By (3.7) and (3.8), we have

$$34 - 2x_1 = \lambda_1^1$$

$$24.25 - (5/4)x_1 = -\lambda_4^2$$

Then $\lambda_4^2 = (5x_1 - 97)/4 < 0$ which contradicts (3.12).

(6) I_5 : Since $x_1 = 0$ and $0 \leq x_2 < 10$, by (3.9), (3.11) and (3.13), we have

$$\lambda_1^i = \lambda_3^i = \lambda_5^i = 0, \quad i = 1, 2$$

By (3.7) and (3.8), we have

$$34 - (8/3)x_2 = -\lambda_2^1$$

$$24.25 - 2x_2 = -\lambda_4^2$$

Then $\lambda_2^1 = (8x_2 - 102)/3 < 0$ which contradicts (3.10).

Hence combining (1) - (6), the solution set of system (3.7) - (3.13) is composed of the point (5,9) and the interval [(9,6), (10,5)].

By Theorem 3.1 and 3.2, the set of nondominated equilibrium points of Γ associated with W is composed of the point (5,9) and the interval [(9,6), (10,5)] (see fig. 1).

REFERENCES

- [1] A. Charnes, W.W. Cooper, Q.L. Wei, and Z.M. Huang, "Fundamental Theorems of Nondominated Solutions Associated with Cones in Normed Linear Spaces," Research Report CCS 575, Center for Cybernetic Studies, The University of Texas at Austin, June, 1987.
- [2] A. Charnes, W.W. Cooper, Q.L. Wei and Z.M. Huang, "Multi-Payoff Constrained N-Person Games Over Cones," Research Report CCS 583, Center for Cybernetic Studies, The University of Texas at Austin, August 1987.
- [3] A. Charnes, W.W. Cooper, Q.L. Wei and Z.M. Huang, "Cone-Ratio Data Envelopment Analysis and Multiobjective Programming," Research Report CCS 559, Center for Cybernetic Studies, The University of Texas at Austin, January, 1987.
- [4] A. Charnes, S.C. Littlechild, M.J.L. Kirby and W.M. Raïke, "Chance-Constrained Models for Transport Pricing and Scheduling Under Competition," Transportation Science, Vol. 14, No. 9, May 1968, p. 513-524.
- [5] P.T. Harker, "Generalized Nash Games and Quasi-Variational Inequalities," Working Paper 86-05-03, Department of Decision Sciences, University of Pennsylvania, Philadelphia, Pennsylvania, May, 1986.
- [6] Z.M. Huang, "The Second Order Conditions of Nondominated Solutions for Generalized Multiobjective Mathematical Programming," Journal of System Science and Mathematical Science, 5 (1985) 3.
- [7] M. Avriel, Nonlinear Programming: Analysis and Methods, Prentice-Hall, Inc., 1976.
- [8] A. Ben-Tal, "Second Order Theory of Extremum Problems," Proceedings of the International Symposium on Extremal Methods and Systems Analysis on the Occasion of Professor A. Charnes' 60th Birthday, 1977.
- [9] P.L. Yu, "Cone Convexity, Cone Extreme Points and Nondominated Solutions in Decision Problems with Multiobjectives," Journal of Optimization Theory and Applications, 14(1974)3.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER CCS 592	2. GOVT ACCESSION NO. 17	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) INTERACTING STRATEGY SETS IN MULTIOBJECTIVE COMPETITION: A DOMINANCE CONE CONSTRAINED GAME SOLUTION		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER CCS 592
7. AUTHOR(s) A. Charnes, Z.M. Huang, J.J. Rousseau, J. Semple		8. CONTRACT OR GRANT NUMBER(s) NSF SES8520806 Army DAKF15-87-C-0110
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Cybernetic Studies University of Texas at Austin Austin, Texas 78712		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS USAREC Ft. Sheridan, IL 60037		12. REPORT DATE April 1988
		13. NUMBER OF PAGES 16
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Dominance Cones, Multi-payoff Cross-Constrained Games, Variational Inequalities		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Models employed in evaluation or stipulation of regulatory policies involving competition and/or differing objectives of competing parties have sometimes been seriously deficient in accounting for interactions between the parties' strategies or in allowing for multiple objectives. Classical game-theoretical models, wherein the strategy set is the topological product of the individual parties strategy sets, do not encompass such situations. The new "dominance cones" method and class of solutions (Charnes, Cooper, Huang and Wei) is herein further extended to such extensions of classical games. This is		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

applied to an example of Harker without requiring his variational and quasi-variational inequalities or point-to-set mappings.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)